

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5:

C07D 401/04, 405/04, 409/04

(11) International Publication Number:

WO 91/07398

A1

(43) International Publication Date:

30 May 1991 (30.05.91)

(21) International Application Number:

PCT/DK90/00304

(22) International Filing Date:

22 November 1990 (22.11.90)

(30) Priority data:

5882/89

22 November 1989 (22.11.89) DK

(71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK).

(72) Inventors: JAKOBSEN, Palle; Langkær Vænge 14, DK-3500 Værløse (DK). SONNEWALD, Ursula; Sildropr-veien 34B, N-7048 Trondheim (NO). TREPPENDAHL, Svend; Frederiksdalsvej 221, DK-2830 Virum (DK).

(74) Agent: LEHMANN & REE; Frederiksberg Allé 26, DK-1820 Frederiksberg C (DK).

(81) Designated States: AT (European patent), AU, BE (European patent), CA, CH (European patent), DE (European patent), DK (European patent), ES (EUROPEAN EUROPEAN EUROPEA tent), FI, FR (European patent), GB (European patent), GR (European patent), HU, IT (European patent), JP, KR, LU (European patent), NL (European patent), NO, SE (European patent).

Published

With international search report.

(54) Title: 4-HETEROARYL PIPERIDINE INTERMEDIATES AND THEIR PREPARATION

(57) Abstract

This invention relates to new compounds of formulae (I, II) in which X is O, S, or NR where R is H or C₁₋₄-alkyl, Z is hydrogen, halogen, trifluoromethyl, C₁₋₈-alkoxy, C₁₋₈-alkyl straight or branched, nitro, C₂₋₈-alkenyl, or a C₁₋₄-alkyl monoor disubstituted amino group, R1 is H or straight or branched C1-8-alkyl. The invention also relates to a method of preparing a compound of formulae (I and II).

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	FI	Finland	ML	Mali
ΑIJ	Australia	FR	France	MN	Mongolia
BB	Barbados	GA	Gabon	MR	Mauritania
BR	Belgium	GB	United Kingdom	MW	Malawi
BP	Burkina Faso	GN	Guinea	NL	Notherlands
BG	Bulgaria	GR	Greece	NO	Norway
BJ	Benin	HU	Hungary	PL	Poland
BR	Brazil	IT	lcaly	RO	Romania
CA	Canada	JP	Japan	SD	Sudan
CF	Central African Republic	KP	Democratic People's Republic	SE	Sweden
CG	Congo		of Korea	SN	Senegal
CH	Switzerland	KR	Republic of Korea	ຮນ	Soviet Union
CI	Côte d'Ivoire	LI	Liochtenstein	TD	Chad
CM	Cameroon	LK	Sri Lanka	TG	Togo
DE	Germany	ш	Luxembourg	US	United States of America
DK	Denmark	MC	Monaco		
ES	Snain	MG	Madagascar		

4-HETEROARYL PIPERIDINE INTERMEDIATES AND THEIR PREPARATION

5

This invention relates to a novel chemical process for preparing new heteroaryl piperidine carbinols and to novel intermediates used in that process.

10

The novel compounds are useful as intermediates in processes leading to pharmacological active substances.

This invention relates to new compounds of formula I

15

25

in which X is O, S, or NR where R is H or C_{1-4} -alkyl, Z is hydrogen, halogen, trifluoromethyl, C_{1-8} -alkoxy, C_{1-8} -alkyl straight or branched, nitro, C_{2-8} -alkenyl, or a C_{1-4} -alkyl mono- or disubstituted amino group, R¹ is H or straight or branched C_{1-8} -alkyl.

The invention also relates to a method of preparing a compound of formula I.

35

30

This method which uses easily accessible commercially available starting materials comprises:

a) preparation of a compound of formula II

 $\begin{array}{c}
z \\
X \\
COOR^2
\end{array}$ II

10

5

wherein X, Z, R and ${\mbox{R}}^1$ have the meaning defined above and ${\mbox{R}}^2$ is ${\mbox{C}}_{1-2}$ -alkyl,

by reacting a compound of formula IV

15

with a compound of the formula V

20

under basic conditions eg. using alkoxide in ethanol, wherein X, Z, R, R^1 and R^2 have the meaning defined above and R^3 being C_{1-2} -alkoxy when R^4 is NHR^1 or R^3 being NHR^1 when R^4 is C_{1-2} -alkoxy,

b) reduction of a compound of formula II wherein X, Z, R, R¹ and R² have the meaning defined above, with metal hydride eg. LiAlH₄ or AlH₃ in ether or THF giving a compound of formula I wherein X, Z, R and R¹ have the meaning defined above, c) reduction of a compound of formula III

10

5

wherein X, Z, R, R^1 and R^2 have the meaning defined above with a metal hydride eg. LiAlH₄ or AlH₃ giving a compound of formula I wherein X, Z, R and R^1 have the meaning defined above;

15


compounds of formula III can be prepared conveniently from Arecoline-type derivatives and metal organic derivatives of heteroaromates using well known procedures,

20

d) compounds of formula I may also be prepared by reacting compounds of formula I wherein Z is H, with reagents causing heteroaromatic substitution using well known procedures,

25

e) compounds of formula I may be prepared by metal hydride reduction of a compound of formula VI



35

wherein X, Z, R, R^1 and R^2 have the meaning defined above,

f) compounds of formula II may be prepared in a one pot reaction using a compound of formula VII wherein X, Z and R have the meaning defined above

5 Z X CHO

as starting material. The reaction is carried out using ethyl acetate as solvent and methoxide or ethoxide as base. After initial reaction between VII and solvent a compound of formula VIII wherein R¹ and R² have the meaning defined above,

15 R²OOCCH₂CONHR¹ VIII

is added resulting in the formation of compounds of formula II wherein X, Z, R, \mathbb{R}^1 and \mathbb{R}^2 have the meaning defined above.

The invention will now be described in further detail with reference to the following examples.

EXAMPLE 1

25

30

3-hydroxymethyl-1-methyl-4-(2-thienyl)-piperidine (1)

3-methoxycarbonyl-1-methyl-4-(2-thienyl)-piperidine (2) was prepared from Arecoline, HBr (52 g), 2-bromothiophene (41 ml) and Mg-turnings (9.9 g) as described by Plati et. al. (J. Org. Chem. 22 (1957) 261). The resulting proudct was purified by distillation giving 15 g cis/trans mixture b.p. 50-120°C / 1.1 mm Hg.

35 30 g of this product was reduced with ${\rm LiAlH_4}$ (5 g) in dry ether (200 ml) by reflux for 30 min. in ${\rm N_2}$ -atmosphere. The well known rinse up procedure gave a hard

oil (21 g) identified as $\frac{3-\text{hydroxymethyl-1-methyl-4-}}{(2-\text{thienyl})-\text{piperidine}(1)}$ by ^1H NMR (CDCl₃), : 6.7 - 7.3 (3H,m); 3.8- 4.5 (1H,broad) 3.5-3.7 (2H,m); 2.8-3.5 (3H,m); 2.2 (3H,s); 1.8-2.8 (5H,m)

5

10

EXAMPLE 2

3-(2-thienyl)-propencyl chloride (3) was prepared by dropwise addition of thionyl chloride (50 ml) to 3-(2-thienyl)-propencic acid (25 g) and subsequent heating to 60°C for 2 hours. Excess of SOCl₂ was evaporated in vacuo, CH₂Cl₂ was added and the reaction mixture was evaporated to dryness yielding 28 g of (3).

N-pentyl-3-(2-thienyl)-propenoic amide (4) was prepared from (3) (28 g) dissolved in dry toluene (200 ml) pentyl amine (25 ml) and triethyl amine (70 ml) was added under cooling (ice bath). Stirring for 1 hour. The precipitate of triethylammonium chloride was removed by filtration, the filtrate was evaporated to dryness and the resulting mass treated with ether giving (4) as colourless crystals (22.8 g). H NMR (CDCl₃),: 0.7-1.1 (3H,m); 1.1-1.7(6H,m); 3.0-3.4 (2H, dist.q), 6.2-8.2 (6H,m).

25

30

35

3-ethoxycarbonyl-1-pentyl-4-(2-thienyl)-piperidine 2,6-dione (5)

Sodium (3 g) was dissolved in abs. ethanol (60 ml), diethyl malonate (20 ml) dissolved in abs. ethanol (30 ml) was added followed by a slurry of (4) (22.8 g) in abs. ethanol (50 ml). Reflux for 4 hours, the ethanol was evaporated, toluene added (100 ml), and the reflux continued overnight. After cooling the formed precipitate was isolated, dissolved in 1M HCl and extracted several times with ether. The combined ether phases were evaporated giving an oil which was purified on silica gel using CH₂Cl₂/CH₃OH 9/1 as eluent. Yield

- 22.6 g of (5) as an oil identified by 1 H NMR (CDCl₃), : 0.7-0.95 (3H,m); 1.0-1.5 (9H,m); 2.9-3.2 (2H,m) 4.15 (2H,q); 3.6-4.15 (4H,m); 6.8-7.2 (3H,m).
- 3-hydroxymethyl-1-pentyl-4-(2-thienyl)-piperidine (6) was prepared from (5) (22.6 g) by reduction with LiAlH₄ (13 g) in dry THF. Reflux for 3 hours under N₂. Using the well known work up procedure gave a hard oil (4 g) identified by ¹H NMR (CDCl₃), : 0.7-1.1 (3H,m); 1.1-1.7 (6H,m); 1.7-2.1 (4H,m); 2.1-3.9 (9H,m); 6.8-7.2 (3H,m).

EXAMPLE 3

- N-pentyl-3-(3-thienyl)-propenoic amide (7) was prepared as described for (4). H NMR (CDCl₃), : 0.7-1.1 (3H,m); 1.1-1.7 (6H,m); 3.2-3.6 (2H, dist. q); 6.1-7.8 (6H,m).
- 3-ethoxycarbonyl-1-pentyl-4-(3-thienyl)-piperidine2,6-dione (8) was prepared as described for (5). 32 g
 (7) gave 41 g of crude (8) which was used without
 further purification. Reduction and work up as described for the preparation of (6) gave 7.7 g of crystalline
 3-hydroxymethyl-1-pentyl-4-(3-thienyl)-piperidine (9).

 M.p. 96.5-97.5°C.

EXAMPLE 4

- 30 (+)-1-butyl-3-ethoxycarbonyl-4-(2-thienyl)-2,6-piperidinedione (10)
- A solution of 2-thiophenecarbaldehyde (22.4 g) in ethyl acetate (20 ml) was added to a slurry of sodium ethanolate (32.6 g) in ethyl acetate (200 ml). The temperature was kept at 10 C and the mixture stirred for one hour. A solution of ethyl N-butylamidomalonate (41.2 g) in

ethyl acetate (40 ml) was slowly added to the mixture whilst keeping the temperature below 5°C . The mixture was stirred for 18 hours at 20°C and neutralized with 25% acetic acid (130 ml). The water phase was discharged and the organic phase extracted twice with saturated sodium chloride solution (2x50 ml). The organic phase was evaporated, the residue dissolved in toluene (200 ml), dried with potassium carbonate and evaporated to give the crude product. Yield 72 g of an oil identified by $^{1}\text{H NMR (CDCl}_{3}$), : 0.7-1.4 (10H,m); 2.7-4.2 (8H,m); 6.7-7.3 (3H,m).

(+)-1-butyl-3-hydroxymethyl-4-(2-thienyl)-piperidine (11)

15

20

25

10

A solution of crude (+)-1-butyl-3-ethoxycarbonyl-4-(2-thienyl)-2,6-piperidindione (72 g) in toluene (100 ml) was added to a slurry of LiAlH₄ (15.2 g) in THF (100 ml) and toluene (50 ml). The temperature was kept below 10° C during the addition. The reaction mixture was stirred for 18 hours and decomposed by careful addition of water (75 ml) keeping the temperature below 10° C. The hydrolyzed mixture was stirred for 1 hour before the precipitated salts was filtered off. The filtrate was evaporated to give an oil (33 g) which was recrystallized from ethyl acetate (50 ml), filtered off and dried to give the title compound (17 g), m.p. 89.7-90.1°C.

30

EXAMPLE 5

(+-)-1-butyl-3-ethoxycarbonyl-4-(5-methyl-2-furanyl)2,6-piperidinedione (12)

was prepared from 5-methyl- 2-furancarbaldehyde (22 g)
35 as described for compound (10). Yield 59.5 g of an oil
(80% pure by HPLC). Identified by ¹H NMR (CDCl₃):
0.7-1.6 (10H, m), 2.2 - 2.4 (3H, d); 2.8 - 4.5 (8H, m);

5.8-7.5 (2H, m).

(+-)-1-butyl-3-hydroxymethyl-4-(5-methyl-2-furanyl)-piperidine (13) was prepared from crude (12) (59 g) by reduction with LiAlH₄, as described for (11). Yield 22 g of (13). M.p. 83.5°C.

EXAMPLE 6

10 (+-)-1-butyl-3-ethoxycarbonyl-4-(1-methyl-2-pyrrolyl)2,6-piperidinedione (14)

was prepared from 1-methyl-2-pyrrolecarbaldehyde (10.2 g) and ethyl N-butylamidomalonate (15.4 g) in ethyl acetate as described for compound (10). The crude product (27 g) was subsequently reduced without further purification as described for compound (11). Yield 14 g of

20 (+-)-1-butyl-3-hydroxymethyl-4-(1-methyl-2-pyrrolyl)piperidine (15) precipitated as the oxalate. ¹H NMR
(CDCl₃): 0.7-1.1 (3H, m); 1.1 - 2.2 (6H, m); 2.5 - 3.8
(10H, m), 3.5 (3H, s), 5.7 - 6.0 (2 H, m); 6.1 - 6.7
(1H, m).

25

EXAMPLE 7

3-Ethoxycarbonyl-1-(2-methylbutyl)-4-(3-thienyl)-2,6-piperidinedione (16)

30

35

was prepared from 3-thiophenecarbaldehyde (20 g) and ethyl N-(2-methylbutylamidomalonate) (35 g) in ethyl acetate as described for compound (10). The crude product (60 g oil) was reduced in THF by means of LiAlH_A as described for compound (11) giving

3-hydroxymethyl-1-(2-methylbutyl)-4-(3-thienyl)-

piperidine (17)

The crude product (23 g) was purified on silica gel using ethyl acetate as eluent.

5

Mass spectrum (M+: 267) degradation in accordance with proposed structure. m.p. 88.8-90.2°C.

EXAMPLE 8

10

The following compounds were prepared exactly as described for (17) using the appropriate substituted thiophenecarbaldehyde and ethyl amidomalonate. The dione intermediate was used without purification.

15

3-hydroxymethyl-1-pentyl-4-(2-thienyl)piperidine (18)

Yield 17.6%; m.p. 107.6°C.

1-butyl-3-hydroxymethyl-4-(2-thienyl)piperidine (19)

Yield 27.2%; m.p. 90.9°C.

1-butyl-3-hydroxymethyl-4-(3-thienyl)piperidine (20)

25

20

Yield 27.9%; m.p. 81.7°C.

3-hydroxymethyl-1-pentyl-4-(3-thienyl)piperidine (21)

Yield 27.5%; m.p. 93.9°C. 30

CLAIMS

 A process for the preparation of a compound of formula I

5

Z CH₂OH

10

15

wherein X is O, S or NR, R is H or C_{1-4} -alkyl, Z is hydrogen, halogen, trifluoromethyl, C_{1-8} -alkoxy, straight or branched C_{1-8} -alkyl, nitro, C_{2-8} -alkenyl, or a C_{1-4} -alkyl mono- or disubstituted amino group,

 R^1 is H, straight or branched C_{1-8} -alkyl;

20

the process comprises a preparation and reduction of a compound of formula II with a metal hydride eg. ${\tt LiAlH}_4$

25

30

35

wherein X, Z, R and R 1 is defined above and R 2 is C_{1-2} -alkyl and the preparation and reduction of a compound of formula III by means of metal hydride eg. LiAlH $_A$

10

15

11.

wherein X, Z, R, R^1 and R^2 have the meaning defined above

2) a compound of formula II

wherein X, Z, R, R^1 , and R^2 have the meaning defined above

20 3) a process for the preparation of a compound of for-

mula II

wherein X, Z, R,
$$R^1$$
 and R^2 have the meaning defined

above, which comprises reacting a compound of formula 30 IV

35 with a compound of formula V

20

30

ð

under basic conditions, wherein

X, Z, R, R^1 and R^2 have the meaning defined above, R^3 being C_{1-2} -alkoxy when R^4 is NHR^1 or R^3 being NHR^1 when R^4 is C_{1-2} -alkoxy.

4) a process for the preparation of a compound of formula II

wherein X, Z, R, ${\rm R}^1$ and ${\rm R}^2$ have the meaning defined above, which comprises reacting of a compound of formula VII

wherein X, Z and R have the meaning defined above, with a compound of formula VIII

R²OOCCH₂CONHR¹ VIII

wherein R^1 and R^2 have the meaning defined above, in ethyl acetate using ethoxide or methoxide as base.

35 5) a compound which is 3-ethoxycarbonyl-1-pentyl-4-(2-thienyl)-piperidine-2,6-dione.

- 6) a compound which is 3-hydroxymethyl-1-methyl-4-(2thienyl)-piperidine.
- 7) a compound which is 3-hydroxymethyl-1-pentyl-4-(3-thienyl)-piperidine.
 - 8) a compound which is 3-ethoxycarbonyl-1-pentyl-4-(3-thienyl)-piperidine-2,6-dione.
- 10 9) a compound which is 3-methoxycarbonyl-1-methyl-4-(2-thienyl)-piperidine.
 - 10) a compound which is 3-hydroxymethyl-1-pentyl-4-(2thienyl)piperidine

11) a compound which is 1-butyl-3-hydroxymethyl-4-(2thienyl)piperidine

- 12) a compound which is 1-butyl-3-hydroxymethyl-4-(5-20 methyl-2-furanyl)piperidine
 - 13) a compound which is 1-butyl-3-hydroxymethyl-4-(3-thienyl)piperidine
- 25 14) a compound which is 3-hydroxymethyl-1-(2-methyl-butyl)-4-(3-thienyl)piperidine
 - 15) a compound which is 1-butyl-3-hydroxymethyl-4-(1-methyl-2-pyrrolyl)piperidine.

30

15

-5

INTERNATIONAL SEARCH REPORT

	International Application No PCI	/DK 90/00304			
1. CLASSIFICATION OF SUBJECT MATTER (if several class					
According to International Patent Classification (IPC) or to both IPC5: C 07 D 401/04, 405/04, 409/04	National Classification and IPC				
II. FIELDS SEARCHED					
Minimum Docum	entation Searched ⁷				
Classification System	Classification Symbols				
IPC5 C 07 D					
	er than Minimum Documentation ts are Included in Fields Searched ⁹				
SE,DK;FI,NO classes as above					
III. DOCUMENTS CONSIDERED TO BE RELEVANTS					
Category * Citation of Document,11 with Indication, where a	propriate, of the relevant passages 12	Relevant to Claim No.13			
X EP, A1, 0223334 (BEECHAM GROUP	PLC)	1-15			
27 May 1987,					
see the whole document					
X EP, A2, 0190496 (BEECHAM GROUP	DI C)	1,3			
13 August 1986,	PLC)	1,3			
see pages 1-19					
Α		2,5-			
		15			
US, A, 4007196 (JORGEN ANDERS) 8 February 1977,	CHRISTENSEN ET AL.)	6-7,9- 15			
see the whole document		}			
.					
	•				
* Special categories of cited documents: 10 "A" document defining the general state of the art which is no considered to be of particular relevance	"T" later document published after or priority date and not in cont cited to understand the princip	the international filing date lict with the application but ble or theory underlying the			
"E" earlier document but published on or after the internation filling date	al "X" document of particular relevan	ice, the claimed invention			
"L" document which may throw doubts on criority claim(s) or which is cited to establish the publication date of another	cannot be considered novel or involve an inventive step	cannot be considered to			
"O" document referring to an oral disclosure, use, exhibition other means.		e or more other such accu-			
"P" document published prior to the international filing date t later than the priority date claimed	in the art.	-			
IV. CERTIFICATION	a cocament member of the semi	s barges caussi			
Date of the Actual Completion of the International Search	Data of Mailing of this International	Search Report			
27th February 1991	1991 -03-	0 6			
International Searching Authority	Signature of Authorized Officer				
	ana Karl				
SWEDISH PATENT OFFICE Göran Karlsson Form PCT/ISA/210 (second sheet) (January 1985)					

International Application No.	PCT/DK90/00304	
FURTHER INFORMATION CONTINUED FROM THE SECOND SHEET		Ì
	•	
	<u>;</u> :	
	:	
	į	
	!	
OBSERVATIONS WHERE CERTAIN CLAIMS WERE FOUND UNSEARCHABLE	····	
This international search report has not been established in respect of certain claims under Article 1	7(2) (a) for the following reasons:	
X Claim numbers 2 decayse they relate to subject matter not required to be searched by t	his Authority, namely:	,
See annexed sheet !	ļ	
os amora succi.		
L. Claim numbers, because they relate to parts of the International application that do not		
ments to such an extent that no meaningful international search can be carried out, specifically	:	
		Ì
		1
Claim numbers, because they are dependent claims and are not drafted in accordance with	th the second and third sentences of	ł
PCT Rule 6.4(a).		
I. ORSERVATIONS WHERE UNITY OF INVENTION IS LACKING 2		
his international Searching Authority found multiple inventions in this international application as fo	ollows:	
As all required additional search fees were timely paid by the applicant, this international search	report covers all searchable claims	
of the international application. As only some of the required additional search fees were timely paid by the applicant, this into		l
those claims of the international application for which fees were paid, specifically claims:	amendum semicu isboir coasis dillà	
	ļ	
No required additional search fees were timely paid by the applicant. Consequently, this interna	tional search report is restricted to	
the invention first mentioned in the claims; it is covered by claim numbers:		
As all searchable claims could be searched without effort justifying an additional fee, the Intern Invite payment of any additional fee.	ational Searching Authority did not	
emark on Protest		
The additional search fees were accompanied by applicant's protest.		
No strategy accompanied the perment of additional search feet		

Claims 2 and 5-15 can not be fully searched and categorized because the structure and the use of the pharmacological active substances are not specified in the description. According to Article 5, the description shall disclose the invention in a manner sufficiently clear and complete for the invention to be carried out by a person skilled in the art.

Form PCT/ISA/210 (supplemental sheet (1)) (October 1981)

ANNEX TO THE INTERNATIONAL SEARCH REPORT ON INTERNATIONAL PATENT APPLICATION NO.PCT/DK 90/00304

This annex lists the patent family members relating to the patent documents cited in the above-mentioned international search report. The members are as contained in the Swedish Patent Office EDP file on 91-01-31 The Swedish Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

Patent document cited in search report	Publication date		family iber(s)	Publication date
EP-A1- 0223334	87-05-27	AU-B- AU-D- JP-A- US-A-	582456 6101286 62039566 4902801	89-03-23 87-02-12 87-02-20 90-02-20
EP-A2- 0190496	86-08-13	AU-D- JP-A-	5111485 61180769	86-06-19 86-08-13
JS-A- 4007196	77-02-08	AT-B- BE-A- CA-A- CH-A- DE-A-C- FR-A-B- GB-A- JP-C- JP-A- JP-B- JP-B- LU-A- NL-A- SE-B-C- US-A- BE-A-	333759 810310 1038390 592059 2404113 2215233 1422263 1268487 1272362 49101385 58174363 59046216 59048826 69264 7401189 401827 3912743 893095	76-12-10 74-05-16 78-09-12 77-10-14 74-08-08 74-08-23 76-01-21 85-06-10 85-07-11 74-09-25 83-10-13 84-11-10 84-11-29 74-04-10 74-08-01 78-05-29 75-10-14 82-08-30



This Page is Inserted by IFW Indexing and Scanning Operations and is not part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:	
☐ BLACK BORDERS	
☐ IMAGE CUT OFF AT TOP, BOTTOM OR SIDES	
☐ FADED TEXT OR DRAWING	
☐ BLURRED OR ILLEGIBLE TEXT OR DRAWING	
☐ SKEWED/SLANTED IMAGES	
☐ COLOR OR BLACK AND WHITE PHOTOGRAPHS	
☐ GRAY SCALE DOCUMENTS	
☐ LINES OR MARKS ON ORIGINAL DOCUMENT	
REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY	

IMAGES ARE BEST AVAILABLE COPY.

OTHER:

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.